

Key Indicators for Arunachal Pradesh from NFHS-3	NFHS-3 (2005-06)	Residence		Education				NFHS-2 (1998-99)	NFHS-1 (1992-93)
		Urban	Rural	No education ⁴	<5 years complete ⁴	5-9 years complete ⁴	10 years complete and above ⁴		
Marriage and Fertility									
1. Women age 20-24 married by age 18 (%)	42.0	43.4	41.3	61.7	(53.3)	40.6	14.6	27.6	43.9
2. Men age 25-29 married by age 21 (%)	30.6	(43.9)	24.7	*	*	31.5	(11.2)	na	na
3. Total fertility rate (children per woman)	3.0	2.5	3.2	4.0	2.8	2.3	1.8	2.5	4.3
4. Women age 15-19 who were already mothers or pregnant at the time of the survey (%)	15.4	18.2	14.1	25.8	21.5	11.3	(2.3)	na	na
5. Median age at first birth for women age 25-49	19.9	20.7	19.7	19.0	18.1	20.7	≥25	20.3	20.1
6. Married women with 2 living children wanting no more children ¹ (%)	73.2	73.4	73.2	62.4	*	73.3	84.1	46.1	31.3
6a. Two sons	85.0	*	(85.0)	*	*	*	*	na	na
6b. One son, one daughter	76.0	(74.4)	76.7	(66.4)	*	(73.9)	(84.5)	na	na
6c. Two daughters	(45.5)	*	*	*	*	*	*	na	na
Family Planning (currently married women, age 15-49)									
Current use									
7. Any method (%)	43.2	47.3	41.6	37.5	43.9	46.3	58.8	35.4	23.6
8. Any modern method (%)	37.3	39.4	36.4	33.4	38.4	38.9	47.8	32.8	19.3
8a. Female sterilization (%)	22.5	19.4	23.7	23.1	30.9	20.0	16.4	20.6	10.3
8b. Male sterilization (%)	0.1	0.0	0.1	0.0	0.0	0.0	0.7	0.1	0.4
8c. IUD (%)	3.4	6.5	2.2	2.2	0.7	4.7	8.3	4.2	4.6
8d. Pill (%)	8.1	8.7	7.9	7.2	4.8	11.3	9.1	7.3	3.2
8e. Condom (%)	2.8	4.5	2.1	0.5	2.0	2.2	13.1	0.7	0.7
Unmet need for family planning									
9. Total unmet need (%)	18.8	18.6	18.9	20.1	19.6	18.7	13.5	26.5	20.4
9a. For spacing (%)	8.3	10.1	7.5	6.8	7.5	12.4	6.8	17.2	12.9
9b. For limiting (%)	10.6	8.5	11.4	13.2	12.1	6.3	6.8	9.3	7.4
Maternal and Child Health									
Maternity care (for births in the last 3 years)									
10. Mothers who had at least 3 antenatal care visits for their last birth (%)	36.4	56.6	28.3	21.5	33.9	50.5	73.5	40.9	31.9
11. Mothers who consumed IFA for 90 days or more when they were pregnant with their last child (%)	11.6	24.3	6.6	4.4	8.1	16.6	36.3	na	na
12. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%) ²	33.6	65.4	21.1	18.3	25.4	47.4	86.3	31.9	22.0
13. Institutional births (%) ²	31.7	64.1	19.0	16.6	28.5	43.2	82.8	31.2	20.1
14. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/other health personnel within 2 days of delivery for their last birth (%) ²	23.3	42.6	15.5	11.4	21.1	28.1	66.7	na	na
Child immunization and vitamin A supplementation²									
15a. Children 12-23 months fully immunized (BCG, measles, and 3 doses each of polio/DPT) (%)	28.4	(51.2)	21.1	19.9	*	(40.8)	*	20.5	22.5
15b. Children 12-23 months who have received BCG (%)	57.7	(65.1)	55.3	49.3	*	(73.7)	*	54.2	46.3
15c. Children 12-23 months who have received 3 doses of polio vaccine (%)	55.8	(62.8)	53.5	49.1	*	(70.6)	*	43.3	38.8
15d. Children 12-23 months who have received 3 doses of DPT vaccine (%)	39.3	(60.5)	32.5	30.8	*	(54.2)	*	41.8	38.8
15e. Children 12-23 months who have received measles vaccine (%)	38.3	(53.5)	33.3	26.2	*	(57.3)	*	33.6	27.5
16. Children age 12-35 months who received a vitamin A dose in last 6 months (%)	18.7	23.6	17.0	11.9	(14.0)	25.5	(42.5)	na	na
Treatment of childhood diseases (children under 3 years)²									
17. Children with diarrhoea in the last 2 weeks who received ORS (%)	33.5	(37.5)	30.8	(27.5)	*	(50.2)	*	40.2	24.1
18. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	38.9	(42.5)	36.5	(29.8)	*	(53.8)	*	52.8	35.6
19. Children with acute respiratory infection or fever in the last 2 weeks taken to a health facility (%)	44.4	(55.8)	38.2	39.3	*	(57.4)	*	na	na
Child Feeding Practices and Nutritional Status of Children^{2,3}									
20. Children under 3 years breastfed within one hour of birth (%)	55.0	53.7	55.5	54.2	62.7	57.7	44.1	49.0	41.4
21. Children age 0-5 months exclusively breastfed (%)	60.0	(55.6)	62.0	(68.5)	*	*	*	na	na
22. Children age 6-9 months receiving solid or semi-solid food and breastmilk (%)	77.6	*	(80.6)	*	*	*	*	na	na
23. Children under 3 years who are stunted (%)	37.0	33.3	37.9	43.5	26.0	33.5	(23.0)	30.3	54.0
24. Children under 3 years who are wasted (%)	17.0	8.7	20.1	24.5	12.9	7.7	(5.0)	10.3	15.5
25. Children under 3 years who are underweight (%)	29.7	15.9	34.9	40.0	20.7	19.3	(11.9)	21.5	32.1
Nutritional Status of Ever-Married Adults (age 15-49)									
26. Women whose Body Mass Index is below normal (%)	15.5	18.6	14.3	19.6	13.0	11.8	7.9	10.7	na
27. Men whose Body Mass Index is below normal (%)	13.6	13.8	13.5	17.6	(17.6)	11.6	9.8	na	na
28. Women who are overweight or obese (%)	10.5	14.2	9.2	6.7	11.4	11.0	24.9	5.1	na
29. Men who are overweight or obese (%)	10.6	10.6	10.5	4.5	(6.9)	10.3	19.5	na	na
Anaemia among Children and Adults									
30. Children age 6-35 months who are anaemic (%)	68.1	72.6	66.3	67.7	(62.8)	71.6	(66.7)	54.5	na
31. Ever-married women age 15-49 who are anaemic (%)	50.6	54.4	49.1	50.9	49.0	49.3	53.9	62.5	na
32. Pregnant women age 15-49 who are anaemic (%)	51.8	*	50.9	(60.5)	*	(51.7)	*	49.2	na
33. Ever-married men age 15-49 who are anaemic (%)	28.5	23.1	30.5	29.2	(35.4)	30.8	21.0	na	na
Knowledge of HIV/AIDS among Ever-Married Adults (age 15-49)									
34. Women who have heard of AIDS (%)	66.2	80.5	60.5	46.3	72.4	88.5	98.0	60.4	16.2
35. Men who have heard of AIDS (%)	74.9	88.1	69.6	44.6	(67.4)	86.0	97.9	na	na
36. Women who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	31.6	47.1	25.6	11.5	32.1	53.3	72.1	na	na
37. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	64.0	76.2	59.1	35.0	(56.9)	70.9	91.6	na	na
Women's Empowerment									
38. Currently married women who usually participate in household decisions (%)	53.5	50.4	54.7	54.7	44.9	55.2	53.9	na	na
39. Ever-married women who have ever experienced spousal violence (%)	38.8	38.0	39.1	45.4	45.2	33.1	15.8	na	na

na : Not available

() Based on 25-49 unweighted cases

* Based on fewer than 25 unweighted cases

¹ Excludes pregnant women.

² Based on the last 2 births in the 3 years before the survey to ever-married women.

³ Based on WHO standard.

⁴ For children education refers to mother's education. Children with missing information on the mother's education are not included in the education columns.