Var Indicators for Chhattiagarh		Resid	lence		Education				
Key Indicators for Chhattisgarh	NEUC 2			No	< 5 years	E Q voors	10 years complete and	NELICO	NEUCA
from NFHS-3	NFHS-3 (2005-06)	Urban	Rural	education <sup>4</sup>	< 5 years complete <sup>4</sup>	complete <sup>4</sup>	above <sup>4</sup>	NFHS-2 (1998-99)	NFHS-1 (1992-93)
Marriage and Fertility	,								
1. Women age 20-24 married by age 18 (%)	55.0	27.1	64.0	82.4	67.1	47.4	10.1	61.3	na
2. Men age 25-29 married by age 21 (%)	44.8	17.5	51.9	(79.5)	(65.3)	50.1	10.5	na	na
3. Total fertility rate (children per woman)	2.6	1.8	2.9	3.2	2.4	2.4	1.9	2.8	na
4. Women age 15-19 who were already mothers or pregnant at the time of the survey (%)	14.6	7.6	16.5	33.9	12.3	9.5	5.4	na	na
5. Median age at first birth for women age 25-49	18.8	20.0	18.5	18.1	18.7	19.5	23.4	17.8	na
6. Married women with 2 living children wanting no more children <sup>1</sup> (%)	75.9	87.3	71.8	67.9	(73.9)	78.3	91.7	53.7	na
6a. Two sons	85.7	94.4	83.3	81.1	*	89.6	(94.5)	72.8	na
6b. One son, one daughter	77.3	88.4	73.1	70.9	*	76.8	90.3	53.5	na
6c. Two daughters	46.8	73.0	(33.3)	(30.0)	*	*	*	22.1	na
Family Planning (currently married women, age 15–49) Current use									
7. Any method (%)	53.2	65.4	49.9	52.6	54.8	49.4	64.4	45.0	na
8. Any modern method (%)	49.1	58.9	46.4	48.6	52.4	46.2	55.6	42.3	na
8a. Female sterilization (%)	40.7	43.9	39.8	42.5	47.5	39.4	27.6	35.1	na
8b. Male sterilization (%)	3.3	1.5	3.8	4.9	2.3	0.7	1.5	3.3	na
8c. IUD (%)	0.8	2.7	0.3	0.3	0.0	0.3	5.5	1.0	na
8d. Pill (%)	1.4	2.9	1.0	0.6	0.3	2.3	4.9	0.8	na
8e. Condom (%)	2.9	7.8	1.6	0.5	2.4	3.5	15.8	2.1	na
Unmet need for family planning									
9. Total unmet need (%)	10.1	9.2	10.3	8.8	9.8	13.4	10.3	13.5	na
9a. For spacing (%)	5.3	4.9	5.4	3.9	3.5	9.4	5.8	8.0	na
9b. For limiting (%)	4.8	4.3	4.9	4.9	6.3	4.0	4.5	5.6	na
Maternal and Child Health									
Maternity care (for births in the last 3 years)	F 4 7	02.2	40.2	44.4	E4 4	62.2	0.6.4	22.2	
10. Mothers who had at least 3 antenatal care visits for their last birth (%) 11. Mothers who consumed IFA for 90 days or more when they were pregnant with their	54.7	82.3	49.2	44.1	51.4	63.3	86.4	33.2	na
last child (%)	21.8	32.1	19.7	13.0	20.3	25.4	56.5	na	na
12. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%) <sup>2</sup>	44.3	74.0	38.5	29.0	58.1	53.0	86.6	32.3	na
13. Institutional births (%) <sup>2</sup>	15.7	58.0	7.5	5.5	9.2	18.0	67.2	13.8	na
14. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/other health									
personnel within 2 days of delivery for their last birth (%) <sup>2</sup>	25.3	63.6	17.6	12.8	23.4	31.0	73.0	na	na
<b>Child immunization and vitamin A supplementation</b> <sup>2</sup> 15a. Children 12-23 months fully immunized (BCG, measles, and 3 doses each of									
polio/DPT) (%)	48.7	74.7	43.1	33.9	*	61.2	(90.9)	21.8	na
15b. Children 12-23 months who have received BCG (%)	84.6	93.3	82.7	79.3	*	87.2	(100.0)	74.3	na
15c. Children 12-23 months who have received 3 doses of polio vaccine (%)	85.1	96.0	82.7	78.6	*	92.6	(95.8)	57.1	na
15d. Children 12-23 months who have received 3 doses of DPT vaccine (%)	62.8	85.3	57.9	51.5	*	70.0	(95.1)	40.9	na
15e. Children 12-23 months who have received measles vaccine (%)	62.5	81.3	58.4	46.5	*	79.6	(100.0)	40.0	na
16. Children age 12-35 months who received a vitamin A dose in last 6 months (%)	14.4	25.7	12.1	8.7	(19.4)	18.3	28.3	na	na
Treatment of childhood diseases (children under 3 years) <sup>2</sup>									
17. Children with diarrhoea in the last 2 weeks who received ORS (%)	42.0	*	(36.4)	(40.7)	*	*	*	29.7	na
18. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	65.3	*	(63.6)	(71.3)	*	*	*	59.3	na
19. Children with acute respiratory infection or fever in the last 2 weeks taken to a health facility (%)	69.5	82.5	66.3	61.7	*	(78.7)	(89.1)	na	na
Child Feeding Practices and Nutritional Status of Children <sup>2, 3</sup>	03.5	02.0	00.5	0		(, 51, )	(6311)		
20. Children under 3 years breastfed within one hour of birth (%)	24.5	27.8	23.8	21.0	20.6	28.0	34.9	13.9	na
21. Children age 0-5 months exclusively breastfed (%)	82.0	(69.2)	84.3	83.5	*	(78.4)	*	na	na
22. Children age 6-9 months receiving solid or semi-solid food and breastmilk (%)	54.5	*	52.1	48.4	*	*	*	na	na
23. Children under 3 years who are stunted (%)	52.6	39.6	55.1	58.9	68.3	46.9	24.0	60.8	na
24. Children under 3 years who are wasted (%)	24.1	24.3	24.1	27.3	33.6	19.3	13.9	24.8	na
25. Children under 3 years who are underweight (%)	47.8	36.1	50.1	54.3	70.5	38.3	22.7	53.2	na
Nutritional Status of Ever-Married Adults (age 15-49)									
26. Women whose Body Mass Index is below normal (%)	41.0	23.5	45.7	45.8	44.2	36.0	19.0	48.1	na
27. Men whose Body Mass Index is below normal (%)	31.8	17.9	35.6	40.1	34.2	34.2	17.6	na	na
28. Women who are overweight or obese (%)	6.7	22.6	2.5	3.0	7.2	8.9	25.2	4.1	na
29. Men who are overweight or obese (%)	6.5	20.1	2.8	2.7	3.3	2.7	18.6	na	na
Anaemia among Children and Adults	00.0	75.6	02.4	0.6.0	00.0	747	70.5	07.7	
30. Children age 6-35 months who are anaemic (%)	80.9 57.6	75.6 50.3	82.1 59.4	86.8 62.1	80.9 53.6	74.7 53.5	70.5	87.7 68.7	na
<ul><li>31. Ever-married women age 15-49 who are anaemic (%)</li><li>32. Pregnant women age 15-49 who are anaemic (%)</li></ul>	57.6 63.1	50.3 65.2	59.4 62.7	62.1 75.5	53.6 (59.3)	53.5 50.4	42.6 (61.0)	68.7 68.3	na na
33. Ever-marrried men age 15-49 who are anaemic (%)	26.4	17.9	28.7	75.5 36.3	(59.3)	24.2	18.9	na	na na
Knowledge of HIV/AIDS among Ever-Married Adults (age 15-49)	۷٠٠٦	17.9	20./	50.5	∠J.J	∠ T.∠	10.5	на	114
34. Women who have heard of AIDS (%)	40.7	78.6	30.7	18.3	48.0	71.8	99.1	19.6	na
35. Men who have heard of AIDS (%)	66.9	98.2	58.5	28.5	54.4	81.1	100.0	na	na
36. Women who know that consistent condom use can reduce the chances of getting		_							
HIV/AIDS (%)	25.9	56.8	17.7	7.5	27.2	48.5	85.1	na	na
37. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	59.6	93.4	50.4	20.9	43.4	74.9	94.0	na	na
Women's Empowerment	33.0	JJ. <del>T</del>	JU.T	20.5	73.7	/ т.У	5-7.0	на	ıα
38. Currently married women who usually participate in household decisions (%)	26.8	35.3	24.5	26.4	30.7	23.0	34.4	na	na
39. Ever-married women who have ever experienced spousal violence (%)	29.9	27.7	30.4	33.7	32.7	26.3	13.0	na	na
				-2.,					

na : Not available

<sup>()</sup> Based on 25-49 unweighted cases

<sup>\*</sup> Based on fewer than 25 unweighted cases

<sup>&</sup>lt;sup>1</sup> Excludes pregnant women.

 $<sup>^{2}</sup>$  Based on the last 2 births in the 3 years before the survey to ever-married women.

<sup>&</sup>lt;sup>3</sup> Based on WHO standard.

<sup>&</sup>lt;sup>4</sup> For children education refers to mother's education. Children with missing information on the mother's education are not included in the education columns.