

Key Indicators for Delhi from NFHS-3	Education					NFHS-2 (1998-99)	NFHS-1 (1992-93)
	NFHS-3 (2005-06)	No education ⁴	<5 years complete ⁴	5-9 years complete ⁴	10 years complete and above ⁴		
Marriage and Fertility							
1. Women age 20-24 married by age 18 (%)	22.7	70.0	*	37.7	6.4	19.8	28.7
2. Men age 25-29 married by age 21 (%)	19.4	(52.4)	*	33.4	7.8	na	na
3. Total fertility rate (children per woman)	2.1	3.5	3.1	2.2	1.7	2.4	3.0
4. Women age 15-19 who were already mothers or pregnant at the time of the survey (%)	5.0	24.6	*	2.9	1.9	na	na
5. Median age at first birth for women age 25-49	21.7	19.4	17.9	20.3	23.7	21.2	20.9
6. Married women with 2 living children wanting no more children ¹ (%)	92.2	80.1	*	87.5	96.3	84.3	78.2
6a. Two sons	95.2	(81.1)	*	96.5	98.3	90.1	86.8
6b. One son, one daughter	94.6	85.6	*	89.1	97.5	89.3	86.0
6c. Two daughters	70.7	*	*	*	(84.1)	52.9	43.8
Family Planning (currently married women, age 15-49)							
Current use							
7. Any method (%)	66.9	59.9	(53.5)	63.9	72.5	63.8	60.3
8. Any modern method (%)	56.5	52.0	(47.1)	54.5	60.2	56.3	54.6
8a. Female sterilization (%)	23.0	32.4	(28.7)	25.9	16.2	26.3	20.0
8b. Male sterilization (%)	0.8	1.4	(1.2)	0.4	0.7	2.3	3.2
8c. IUD (%)	5.0	2.0	(2.9)	5.3	6.5	6.2	7.8
8d. Pill (%)	4.5	4.6	(5.7)	4.6	4.4	4.0	2.9
8e. Condom (%)	22.9	11.2	(8.7)	18.3	31.9	17.5	20.5
Unmet need for family planning							
9. Total unmet need (%)	7.8	10.8	(25.3)	10.8	4.2	13.4	15.4
9a. For spacing (%)	3.3	3.9	(9.2)	5.0	2.1	5.9	7.6
9b. For limiting (%)	4.5	6.9	(16.1)	5.8	2.1	7.5	7.9
Maternal and Child Health							
Maternity care (for births in the last 3 years)							
10. Mothers who had at least 3 antenatal care visits for their last birth (%)	74.4	51.9	*	69.2	93.0	68.9	72.2
11. Mothers who consumed IFA for 90 days or more when they were pregnant with their last child (%)	38.3	13.6	*	26.3	62.8	na	na
12. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%) ²	65.1	32.8	*	59.1	93.2	65.9	53.8
13. Institutional births (%) ²	60.7	26.8	*	54.0	91.0	59.1	45.3
14. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/other health personnel within 2 days of delivery for their last birth (%) ²	50.4	24.5	*	34.5	78.0	na	na
Child immunization and vitamin A supplementation²							
15a. Children 12-23 months fully immunized (BCG, measles, and 3 doses each of polio/DPT) (%)	63.2	44.2	*	(67.2)	76.2	69.8	57.8
15b. Children 12-23 months who have received BCG (%)	87.0	68.1	*	(90.2)	100.0	92.0	90.1
15c. Children 12-23 months who have received 3 doses of polio vaccine (%)	79.1	66.2	*	(85.8)	86.0	81.0	75.6
15d. Children 12-23 months who have received 3 doses of DPT vaccine (%)	71.7	49.6	*	(74.0)	86.5	79.9	71.6
15e. Children 12-23 months who have received measles vaccine (%)	78.2	55.4	*	(82.1)	93.3	77.5	69.6
16. Children age 12-35 months who received a vitamin A dose in last 6 months (%)	20.3	12.6	*	18.5	26.4	na	na
Treatment of childhood diseases (children under 3 years)²							
17. Children with diarrhoea in the last 2 weeks who received ORS (%)	34.4	(27.5)	*	*	*	39.1	18.9
18. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	75.1	(73.6)	*	*	*	80.1	66.2
19. Children with acute respiratory infection or fever in the last 2 weeks taken to a health facility (%)	93.1	(91.9)	*	(90.9)	(95.5)	na	na
Child Feeding Practices and Nutritional Status of Children^{2,3}							
20. Children under 3 years breastfed within one hour of birth (%)	19.3	17.8	*	13.7	24.2	23.8	6.3
21. Children age 0-5 months exclusively breastfed (%)	34.5	(36.4)	*	(31.8)	(31.9)	na	na
22. Children age 6-9 months receiving solid or semi-solid food and breastmilk (%)	59.8	(61.4)	*	*	(52.4)	na	na
23. Children under 3 years who are stunted (%)	43.2	54.2	*	49.5	31.1	43.2	46.9
24. Children under 3 years who are wasted (%)	17.2	19.6	*	13.5	18.4	15.7	15.9
25. Children under 3 years who are underweight (%)	24.9	34.8	*	26.1	17.5	29.9	36.2
Nutritional Status of Ever-Married Adults (age 15-49)							
26. Women whose Body Mass Index is below normal (%)	10.6	15.4	(14.5)	14.6	6.0	12.0	na
27. Men whose Body Mass Index is below normal (%)	10.4	20.7	*	13.3	6.2	na	na
28. Women who are overweight or obese (%)	32.9	24.2	(10.7)	29.0	40.2	33.8	na
29. Men who are overweight or obese (%)	24.0	17.4	*	16.4	29.7	na	na
Anaemia among Children and Adults							
30. Children age 6-35 months who are anaemic (%)	63.2	71.4	*	67.6	53.9	69.0	na
31. Ever-married women age 15-49 who are anaemic (%)	43.3	48.9	(42.3)	43.2	40.3	40.5	na
32. Pregnant women age 15-49 who are anaemic (%)	29.9	(36.2)	*	(30.9)	(22.8)	34.7	na
33. Ever-married men age 15-49 who are anaemic (%)	18.9	28.5	*	22.5	14.9	na	na
Knowledge of HIV/AIDS among Ever-Married Adults (age 15-49)							
34. Women who have heard of AIDS (%)	88.1	65.4	86.1	91.8	99.4	79.2	35.8
35. Men who have heard of AIDS (%)	97.0	84.4	*	97.5	99.7	na	na
36. Women who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	77.0	47.3	58.2	76.2	95.0	na	na
37. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	92.4	69.4	*	93.6	97.5	na	na
Women's Empowerment							
38. Currently married women who usually participate in household decisions (%)	52.0	50.4	(36.2)	50.1	54.3	na	na
39. Ever-married women who have ever experienced spousal violence (%)	16.3	28.9	(28.9)	20.7	6.5	na	na

na : Not available

nc : Not calculated because there are no cases

() Based on 25-49 unweighted cases

* Based on fewer than 25 unweighted cases

¹ Excludes pregnant women.

² Based on the last 2 births in the 3 years before the survey to ever-married women.

³ Based on WHO standard.

⁴ For children education refers to mother's education. Children with missing information on the mother's education are not included in the education columns.