

Key Indicators for Goa from NFHS-3	Residence			Education				NFHS-2 (1998-99)	NFHS-1 (1992-93)
	NFHS-3 (2005-06)	Urban	Rural	No education ⁴	<5 years complete ⁴	5-9 years complete ⁴	10 years complete and above ⁴		
Marriage and Fertility									
1. Women age 20-24 married by age 18 (%)	12.1	15.0	8.2	(62.8)	*	12.1	4.8	10.1	7.2
2. Men age 25-29 married by age 21 (%)	7.2	7.9	6.5	*	*	8.0	1.5	na	na
3. Total fertility rate (children per woman)	1.8	1.8	1.8	2.9	1.6	1.9	1.8	1.8	1.9
4. Women age 15-19 who were already mothers or pregnant at the time of the survey (%)	3.6	5.4	1.2	*	*	4.2	1.2	na	na
5. Median age at first birth for women age 25-49	≥25	≥25	≥25	20.7	23.7	≥25	≥25	24.1	23.7
6. Married women with 2 living children wanting no more children ¹ (%)	83.0	83.5	82.3	77.5	78.5	83.0	84.8	75.3	70.3
6a. Two sons	86.3	84.2	89.1	(79.5)	*	83.7	89.1	77.3	76.0
6b. One son, one daughter	84.8	84.9	84.7	(79.5)	(73.2)	86.7	86.3	86.4	76.4
6c. Two daughters	71.7	77.4	63.8	*	*	(71.6)	74.6	52.1	48.9
Family Planning (currently married women, age 15-49)									
Current use									
7. Any method (%)	48.2	51.3	44.1	56.4	50.9	42.9	48.2	47.5	47.8
8. Any modern method (%)	37.2	38.6	35.2	52.3	45.4	35.1	31.3	35.9	37.9
8a. Female sterilization (%)	25.8	24.5	27.4	49.7	42.5	26.8	12.7	27.8	29.5
8b. Male sterilization (%)	0.1	0.1	0.0	0.0	0.0	0.0	0.1	0.4	1.0
8c. IUD (%)	2.2	2.6	1.8	0.5	1.0	2.1	3.2	1.9	2.7
8d. Pill (%)	1.5	1.9	1.0	0.6	0.0	1.1	2.5	0.9	0.7
8e. Condom (%)	7.5	9.3	5.0	1.5	1.3	5.1	12.6	4.9	3.9
Unmet need for family planning									
9. Total unmet need (%)	13.1	12.0	14.6	5.2	10.2	14.9	15.4	17.1	15.7
9a. For spacing (%)	7.4	6.7	8.3	3.6	4.1	9.1	8.3	7.3	7.8
9b. For limiting (%)	5.7	5.3	6.2	1.6	6.1	5.8	7.1	9.8	7.9
Maternal and Child Health									
Maternity care (for births in the last 3 years)									
10. Mothers who had at least 3 antenatal care visits for their last birth (%)	95.0	96.4	93.2	79.3	(86.0)	96.1	99.4	96.3	89.8
11. Mothers who consumed IFA for 90 days or more when they were pregnant with their last child (%)	72.0	78.9	63.3	48.0	(62.2)	71.2	80.0	na	na
12. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%) ²	94.3	94.6	93.8	77.2	(78.9)	95.5	100.0	90.8	89.2
13. Institutional births (%) ²	92.6	92.6	92.4	67.5	(78.9)	94.9	99.7	90.9	87.8
14. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/other health personnel within 2 days of delivery for their last birth (%) ²	78.4	80.4	75.8	69.8	(68.7)	80.6	80.2	na	na
Child immunization and vitamin A supplementation²									
15a. Children 12-23 months fully immunized (BCG, measles, and 3 doses each of polio/DPT) (%)	78.6	83.5	73.0	(52.4)	*	76.9	87.8	82.6	74.9
15b. Children 12-23 months who have received BCG (%)	96.8	97.9	95.5	(89.9)	*	97.2	99.1	99.2	93.5
15c. Children 12-23 months who have received 3 doses of polio vaccine (%)	87.2	88.7	85.6	(70.9)	*	87.5	92.2	95.8	87.8
15d. Children 12-23 months who have received 3 doses of DPT vaccine (%)	87.5	90.7	83.8	(72.7)	*	87.8	91.9	93.4	86.7
15e. Children 12-23 months who have received measles vaccine (%)	91.2	93.8	88.3	(66.6)	*	91.5	99.1	84.3	77.8
16. Children age 12-35 months who received a vitamin A dose in last 6 months (%)	41.9	39.3	45.2	37.6	*	34.9	49.2	na	na
Treatment of childhood diseases (children under 3 years)²									
17. Children with diarrhoea in the last 2 weeks who received ORS (%)	51.6	*	(62.1)	*	*	*	*	55.6	26.9
18. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	71.5	*	(79.3)	*	*	*	*	65.4	67.9
19. Children with acute respiratory infection or fever in the last 2 weeks taken to a health facility (%)	83.7	88.5	79.0	*	*	88.5	81.4	na	na
Child Feeding Practices and Nutritional Status of Children^{2,3}									
20. Children under 3 years breastfed within one hour of birth (%)	59.7	64.7	53.4	65.9	(74.3)	58.9	57.2	34.4	29.1
21. Children age 0-5 months exclusively breastfed (%)	17.7	(19.0)	(15.8)	*	*	*	(13.9)	na	na
22. Children age 6-9 months receiving solid or semi-solid food and breastmilk (%)	69.8	(62.2)	(78.6)	*	*	(77.5)	(60.3)	na	na
23. Children under 3 years who are stunted (%)	25.9	22.8	29.3	47.7	*	27.4	16.4	21.7	34.9
24. Children under 3 years who are wasted (%)	12.8	9.6	17.2	21.0	*	14.1	10.1	16.5	17.5
25. Children under 3 years who are underweight (%)	21.3	15.1	28.9	38.5	*	25.3	12.2	21.3	29.3
Nutritional Status of Ever-Married Adults (age 15-49)									
26. Women whose Body Mass Index is below normal (%)	20.5	16.1	26.1	31.6	25.2	22.5	12.9	27.1	na
27. Men whose Body Mass Index is below normal (%)	16.8	14.5	20.0	24.5	33.3	19.3	8.1	na	na
28. Women who are overweight or obese (%)	27.0	32.1	20.4	17.2	15.1	26.1	35.0	21.2	na
29. Men who are overweight or obese (%)	20.8	23.2	17.3	9.8	3.0	19.9	28.7	na	na
Anaemia among Children and Adults									
30. Children age 6-35 months who are anaemic (%)	49.2	49.4	49.2	57.8	*	53.7	41.7	53.4	na
31. Ever-married women age 15-49 who are anaemic (%)	39.1	38.9	38.8	50.0	40.3	39.8	32.9	36.4	na
32. Pregnant women age 15-49 who are anaemic (%)	36.9	43.3	(26.1)	*	*	(45.6)	23.5	35.0	na
33. Ever-married men age 15-49 who are anaemic (%)	12.9	13.9	11.4	26.6	18.4	12.4	8.3	na	na
Knowledge of HIV/AIDS among Ever-Married Adults (age 15-49)									
34. Women who have heard of AIDS (%)	83.3	85.0	81.0	47.8	70.7	88.1	97.9	76.3	41.7
35. Men who have heard of AIDS (%)	92.2	93.9	89.7	62.8	82.4	95.9	98.2	na	na
36. Women who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	48.8	51.6	45.1	18.8	29.7	47.7	66.7	na	na
37. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	57.3	63.6	48.3	19.1	39.0	59.1	69.7	na	na
Women's Empowerment									
38. Currently married women who usually participate in household decisions (%)	47.0	43.9	51.3	38.4	42.7	40.8	55.4	na	na
39. Ever-married women who have ever experienced spousal violence (%)	16.8	16.4	17.2	35.9	20.1	15.8	8.8	na	na

na : Not available

nc : Not calculated because there are no cases

() Based on 25-49 unweighted cases

* Based on fewer than 25 unweighted cases

¹ Excludes pregnant women.

² Based on the last 2 births in the 3 years before the survey to ever-married women.

³ Based on WHO standard.

⁴ For children education refers to mother's education. Children with missing information on the mother's education are not included in the education columns.