

Key Indicators for Manipur from NFHS-3	NFHS-3 (2005-06)	Residence		Education				NFHS-2 (1998-99)	NFHS-1 (1992-93)
		Urban	Rural	No education <sup>4</sup>	<5 years complete <sup>4</sup>	5-9 years complete <sup>4</sup>	10 years		
							complete and above <sup>4</sup>		
<b>Marriage and Fertility</b>									
1. Women age 20-24 married by age 18 (%)	12.9	11.7	13.5	29.5	(23.9)	16.7	2.5	9.9	14.3
2. Men age 25-29 married by age 21 (%)	11.6	9.2	12.9	*	(17.8)	18.6	5.5	na	na
3. Total fertility rate (children per woman)	2.8	2.4	3.1	3.7	3.4	2.9	2.5	3.0	2.8
4. Women age 15-19 who were already mothers or pregnant at the time of the survey (%)	7.3	4.2	8.6	25.4	11.9	5.6	3.8	na	na
5. Median age at first birth for women age 25-49	23.7	≥25	22.9	21.0	21.7	22.3	≥25	23.1	22.4
6. Married women with 2 living children wanting no more children <sup>1</sup> (%)	66.1	72.0	63.0	64.4	53.2	65.1	70.6	45.6	39.5
6a. Two sons	68.8	81.0	62.1	(65.6)	*	(71.9)	75.1	na	na
6b. One son, one daughter	76.2	76.4	76.0	74.9	*	75.1	79.1	na	na
6c. Two daughters	24.3	(35.0)	19.6	*	*	(17.8)	(28.2)	na	na
<b>Family Planning (currently married women, age 15-49)</b>									
<b>Current use</b>									
7. Any method (%)	48.7	54.5	46.0	43.2	48.6	50.8	51.9	38.7	34.9
8. Any modern method (%)	23.6	24.8	23.0	23.9	22.1	25.1	22.1	25.9	24.1
8a. Female sterilization (%)	8.2	8.9	7.9	11.9	11.0	6.4	5.6	14.4	10.9
8b. Male sterilization (%)	0.5	0.6	0.4	0.3	0.5	0.6	0.6	1.1	2.9
8c. IUD (%)	5.3	4.0	5.9	4.1	4.4	7.0	5.1	6.8	6.7
8d. Pill (%)	5.3	4.7	5.6	5.5	4.8	6.8	3.8	2.2	2.4
8e. Condom (%)	4.1	6.4	3.1	2.0	1.5	4.1	7.0	1.3	1.2
<b>Unmet need for family planning</b>									
9. Total unmet need (%)	12.4	11.9	12.6	12.6	10.4	12.2	13.0	23.6	21.7
9a. For spacing (%)	5.0	5.9	4.5	3.9	2.8	5.8	5.8	13.6	11.7
9b. For limiting (%)	7.4	6.0	8.1	8.7	7.6	6.4	7.2	10.0	10.0
<b>Maternal and Child Health</b>									
<b>Maternity care (for births in the last 3 years)</b>									
10. Mothers who had at least 3 antenatal care visits for their last birth (%)	70.1	85.9	64.0	45.7	52.4	75.5	86.6	54.7	41.3
11. Mothers who consumed IFA for 90 days or more when they were pregnant with their last child (%)	13.9	21.8	10.8	4.2	8.9	11.7	24.3	na	na
12. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%) <sup>2</sup>	61.7	85.2	52.8	34.5	43.3	65.4	82.6	53.9	39.9
13. Institutional births (%) <sup>2</sup>	49.3	71.2	40.9	22.5	32.9	49.3	73.1	34.5	23.0
14. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/other health personnel within 2 days of delivery for their last birth (%) <sup>2</sup>	49.1	70.1	40.9	24.4	31.4	47.5	73.2	na	na
<b>Child immunization and vitamin A supplementation<sup>2</sup></b>									
15a. Children 12-23 months fully immunized (BCG, measles, and 3 doses each of polio/DPT) (%)	46.8	59.6	42.8	25.0	*	47.7	62.4	42.3	29.1
15b. Children 12-23 months who have received BCG (%)	80.0	89.5	77.0	59.1	*	79.3	95.8	71.0	63.8
15c. Children 12-23 months who have received 3 doses of polio vaccine (%)	77.5	83.3	75.7	56.0	*	83.4	88.6	62.5	39.4
15d. Children 12-23 months who have received 3 doses of DPT vaccine (%)	61.2	75.4	56.8	36.7	*	64.5	77.8	59.1	43.3
15e. Children 12-23 months who have received measles vaccine (%)	52.8	64.9	49.1	32.1	*	54.3	68.2	45.8	37.0
16. Children age 12-35 months who received a vitamin A dose in last 6 months (%)	15.2	21.4	12.8	9.8	(11.5)	13.4	21.7	na	na
<b>Treatment of childhood diseases (children under 3 years)<sup>2</sup></b>									
17. Children with diarrhoea in the last 2 weeks who received ORS (%)	36.8	33.3	38.5	34.2	22.2	44.4	33.7	50.7	57.4
18. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	42.2	52.4	37.2	48.8	11.1	35.9	51.6	44.8	42.6
19. Children with acute respiratory infection or fever in the last 2 weeks taken to a health facility (%)	49.1	57.5	45.0	(33.8)	*	52.8	55.2	na	na
<b>Child Feeding Practices and Nutritional Status of Children<sup>2,3</sup></b>									
20. Children under 3 years breastfed within one hour of birth (%)	57.2	58.4	56.8	54.9	49.4	59.1	59.0	27.0	13.2
21. Children age 0-5 months exclusively breastfed (%)	61.7	71.6	57.9	(57.8)	*	66.2	60.0	na	na
22. Children age 6-9 months receiving solid or semi-solid food and breastmilk (%)	78.1	83.3	76.2	(72.7)	*	75.9	90.2	na	na
23. Children under 3 years who are stunted (%)	29.0	26.1	30.0	37.7	35.9	26.9	23.6	38.5	32.0
24. Children under 3 years who are wasted (%)	10.8	10.5	10.8	12.5	16.3	12.6	6.2	9.7	9.9
25. Children under 3 years who are underweight (%)	19.5	17.1	20.3	27.9	25.4	20.4	11.5	20.1	19.1
<b>Nutritional Status of Ever-Married Adults (age 15-49)</b>									
26. Women whose Body Mass Index is below normal (%)	13.9	10.1	15.5	17.5	15.4	14.8	8.5	18.8	na
27. Men whose Body Mass Index is below normal (%)	12.2	10.6	13.0	15.7	23.8	13.9	8.6	na	na
28. Women who are overweight or obese (%)	17.6	27.7	13.1	13.9	12.7	15.1	25.8	10.8	na
29. Men who are overweight or obese (%)	13.4	21.9	9.5	4.7	8.0	7.2	20.3	na	na
<b>Anaemia among Children and Adults</b>									
30. Children age 6-35 months who are anaemic (%)	52.8	53.6	52.5	47.0	59.9	57.5	50.4	45.2	na
31. Ever-married women age 15-49 who are anaemic (%)	39.3	40.6	38.7	38.2	38.9	39.9	40.0	28.9	na
32. Pregnant women age 15-49 who are anaemic (%)	36.4	37.1	36.1	36.8	*	41.0	32.5	36.7	na
33. Ever-married men age 15-49 who are anaemic (%)	14.1	13.8	14.2	19.5	16.6	15.9	11.4	na	na
<b>Knowledge of HIV/AIDS among Ever-Married Adults (age 15-49)</b>									
34. Women who have heard of AIDS (%)	98.5	99.4	98.1	95.4	99.6	99.8	100.0	92.9	72.5
35. Men who have heard of AIDS (%)	99.3	99.5	99.1	93.9	98.5	99.7	100.0	na	na
36. Women who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	80.7	90.3	76.2	64.4	74.4	86.2	93.8	na	na
37. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	93.9	96.1	92.7	78.7	86.7	94.2	97.4	na	na
<b>Women's Empowerment</b>									
38. Currently married women who usually participate in household decisions (%)	69.4	69.2	69.6	78.5	69.5	63.8	66.4	na	na
39. Ever-married women who have ever experienced spousal violence (%)	43.8	44.4	43.5	46.9	52.7	44.4	36.8	na	na

na : Not available

( ) Based on 25-49 unweighted cases

\* Based on fewer than 25 unweighted cases

<sup>1</sup> Excludes pregnant women.

<sup>2</sup> Based on the last 2 births in the 3 years before the survey to ever-married women.

<sup>3</sup> Based on WHO standard.

<sup>4</sup> For children education refers to mother's education. Children with missing information on the mother's education are not included in the education columns.