

Key Indicators for Nagaland from NFHS-3	Residence			Education				NFHS-2 (1998-99)	NFHS-1 (1992-93)
	NFHS-3 (2005-06)	Urban	Rural	No education <sup>4</sup>	<5 years complete <sup>4</sup>	5-9 years complete <sup>4</sup>	10 years		
							complete and above <sup>4</sup>		
<b>Marriage and Fertility</b>									
1. Women age 20-24 married by age 18 (%)	21.4	15.2	24.0	50.4	26.1	20.3	4.6	22.9	16.4
2. Men age 25-29 married by age 21 (%)	18.2	12.1	21.0	28.8	24.0	21.0	9.3	na	na
3. Total fertility rate (children per woman)	3.7	2.7	4.2	5.4	3.8	3.5	2.2	3.8	3.3
4. Women age 15-19 who were already mothers or pregnant at the time of the survey (%)	7.5	8.9	6.8	20.2	10.4	5.7	3.7	na	na
5. Median age at first birth for women age 25-49	21.8	22.6	21.6	20.3	20.9	21.6	≥25	21.2	21.2
6. Married women with 2 living children wanting no more children <sup>1</sup> (%)	58.4	69.6	51.8	54.7	(40.7)	58.5	68.8	47.8	34.7
6a. Two sons	62.2	67.9	(57.5)	*	*	48.7	(77.6)	na	na
6b. One son, one daughter	60.7	75.0	53.5	52.4	(38.5)	67.6	68.3	na	na
6c. Two daughters	46.7	57.7	(40.5)	*	*	(48.4)	*	na	na
<b>Family Planning (currently married women, age 15-49)</b>									
<b>Current use</b>									
7. Any method (%)	29.7	41.9	24.8	17.0	25.5	34.1	43.0	30.3	13.0
8. Any modern method (%)	22.5	31.8	18.8	12.4	19.0	26.2	33.0	24.2	13.0
8a. Female sterilization (%)	9.9	14.8	8.0	4.5	11.5	12.2	12.2	12.3	6.3
8b. Male sterilization (%)	0.0	0.2	0.0	0.0	0.0	0.1	0.1	0.0	0.1
8c. IUD (%)	5.2	4.6	5.5	2.9	2.0	7.1	7.1	7.7	2.0
8d. Pill (%)	4.7	6.5	3.9	4.3	3.6	4.9	5.6	2.5	2.1
8e. Condom (%)	2.6	5.6	1.5	0.8	1.8	2.0	7.7	1.8	2.1
<b>Unmet need for family planning</b>									
9. Total unmet need (%)	26.1	20.2	28.5	29.9	28.9	25.5	19.5	30.2	26.7
9a. For spacing (%)	10.0	7.8	10.9	9.1	12.5	10.2	9.1	18.3	12.9
9b. For limiting (%)	16.1	12.4	17.6	20.8	16.4	15.3	10.4	11.9	13.8
<b>Maternal and Child Health</b>									
<b>Maternity care (for births in the last 3 years)</b>									
10. Mothers who had at least 3 antenatal care visits for their last birth (%)	31.6	57.9	23.9	17.6	23.6	33.9	66.4	21.9	15.0
11. Mothers who consumed IFA for 90 days or more when they were pregnant with their last child (%)	3.5	9.0	1.8	1.5	1.8	2.4	12.9	na	na
12. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%) <sup>2</sup>	25.9	54.3	17.9	10.2	17.3	29.2	60.8	32.8	18.9
13. Institutional births (%) <sup>2</sup>	12.2	32.0	6.6	4.9	8.0	12.6	31.8	12.1	6.1
14. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/other health personnel within 2 days of delivery for their last birth (%) <sup>2</sup>	11.3	30.5	5.7	5.0	6.9	11.8	29.3	na	na
<b>Child immunization and vitamin A supplementation<sup>2</sup></b>									
15a. Children 12-23 months fully immunized (BCG, measles, and 3 doses each of polio/DPT) (%)	21.0	36.0	17.1	11.5	6.1	22.8	49.1	14.1	3.8
15b. Children 12-23 months who have received BCG (%)	46.3	67.7	40.9	29.5	30.1	51.8	79.6	46.1	19.4
15c. Children 12-23 months who have received 3 doses of polio vaccine (%)	46.2	73.2	39.3	31.0	22.3	51.7	83.9	41.8	16.3
15d. Children 12-23 months who have received 3 doses of DPT vaccine (%)	28.7	48.2	23.7	16.7	10.5	32.6	59.1	29.6	12.5
15e. Children 12-23 months who have received measles vaccine (%)	27.3	47.6	22.2	14.1	14.0	29.7	60.0	19.6	10.0
16. Children age 12-35 months who received a vitamin A dose in last 6 months (%)	9.1	17.8	6.7	4.8	6.4	8.3	22.9	na	na
<b>Treatment of childhood diseases (children under 3 years)<sup>2</sup></b>									
17. Children with diarrhoea in the last 2 weeks who received ORS (%)	17.1	(32.7)	(10.6)	*	*	(13.9)	*	29.7	16.7
18. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	19.5	(30.6)	(14.9)	*	*	(19.4)	*	23.3	11.1
19. Children with acute respiratory infection or fever in the last 2 weeks taken to a health facility (%)	23.2	31.3	19.8	12.0	*	30.4	(26.0)	na	na
<b>Child Feeding Practices and Nutritional Status of Children<sup>2,3</sup></b>									
20. Children under 3 years breastfed within one hour of birth (%)	51.5	49.3	52.1	54.8	57.2	46.2	53.5	24.5	66.0
21. Children age 0-5 months exclusively breastfed (%)	29.2	25.4	30.3	(35.1)	*	28.4	(25.6)	na	na
22. Children age 6-9 months receiving solid or semi-solid food and breastmilk (%)	71.0	76.7	69.2	(65.6)	*	71.7	*	na	na
23. Children under 3 years who are stunted (%)	34.1	24.2	36.5	40.3	36.0	31.1	25.8	38.7	32.7
24. Children under 3 years who are wasted (%)	15.8	13.3	16.4	22.0	14.2	12.9	10.6	13.6	12.2
25. Children under 3 years who are underweight (%)	23.7	18.1	25.0	29.7	24.7	20.7	16.6	18.8	18.7
<b>Nutritional Status of Ever-Married Adults (age 15-49)</b>									
26. Women whose Body Mass Index is below normal (%)	15.9	13.1	16.9	19.0	13.7	16.0	11.9	18.4	na
27. Men whose Body Mass Index is below normal (%)	10.8	9.6	11.3	16.7	14.6	8.8	7.7	na	na
28. Women who are overweight or obese (%)	8.9	17.5	5.6	5.8	8.1	9.2	14.2	8.2	na
29. Men who are overweight or obese (%)	8.4	16.8	5.0	3.1	3.1	7.5	16.2	na	na
<b>Anaemia among Children and Adults</b>									
30. Children age 6-35 months who are anaemic (%)	na	na	na	na	na	na	na	43.7	na
31. Ever-married women age 15-49 who are anaemic (%)	na	na	na	na	na	na	na	38.4	na
32. Pregnant women age 15-49 who are anaemic (%)	na	na	na	na	na	na	na	38.2	na
33. Ever-married men age 15-49 who are anaemic (%)	na	na	na	na	na	na	na	na	na
<b>Knowledge of HIV/AIDS among Ever-Married Adults (age 15-49)</b>									
34. Women who have heard of AIDS (%)	80.9	89.4	77.4	61.8	70.7	89.7	98.8	72.4	40.9
35. Men who have heard of AIDS (%)	90.8	94.1	89.4	74.7	86.5	94.3	99.4	na	na
36. Women who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	39.7	54.1	34.0	19.8	32.9	46.6	61.4	na	na
37. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	68.6	75.4	65.6	49.6	57.2	72.0	82.8	na	na
<b>Women's Empowerment</b>									
38. Currently married women who usually participate in household decisions (%)	73.1	70.5	74.1	65.2	72.4	75.4	81.0	na	na
39. Ever-married women who have ever experienced spousal violence (%)	15.3	15.6	15.2	19.6	19.9	13.1	10.3	na	na

na : Not available

nc : Not calculated because there are no cases

( ) Based on 25-49 unweighted cases

\* Based on fewer than 25 unweighted cases

<sup>1</sup> Excludes pregnant women.

<sup>2</sup> Based on the last 2 births in the 3 years before the survey to ever-married women.

<sup>3</sup> Based on WHO standard.

<sup>4</sup> For children education refers to mother's education. Children with missing information on the mother's education are not included in the education columns.