

Key Indicators for Rajasthan from NFHS-3	Residence		Education				NFHS-2 (1998-99)	NFHS-1 (1992-93)	
	NFHS-3 (2005-06)	Urban	Rural	No education ⁴	<5 years complete ⁴	5-9 years complete ⁴			10 years
							complete and above ⁴		
Marriage and Fertility									
1. Women age 20-24 married by age 18 (%)	65.2	38.3	76.0	82.4	(69.5)	64.5	11.3	68.3	69.5
2. Men age 25-29 married by age 21 (%)	57.0	35.4	66.2	(73.8)	*	55.7	44.7	na	na
3. Total fertility rate (children per woman)	3.2	2.2	3.6	3.7	3.0	2.5	1.8	3.8	3.6
4. Women age 15-19 who were already mothers or pregnant at the time of the survey (%)	16.0	8.1	19.0	22.8	19.9	13.7	3.0	na	na
5. Median age at first birth for women age 25-49	19.6	20.1	19.4	19.3	19.2	19.6	23.7	19.4	19.7
6. Married women with 2 living children wanting no more children ¹ (%)	74.8	86.0	67.5	65.9	*	78.1	93.6	58.3	44.2
6a. Two sons	83.9	92.1	79.2	76.5	*	(92.4)	(95.6)	74.8	57.4
6b. One son, one daughter	77.7	90.5	68.9	66.2	*	83.5	95.7	63.1	53.0
6c. Two daughters	33.1	(51.4)	(18.2)	(26.0)	*	*	*	12.5	6.7
Family Planning (currently married women, age 15-49)									
Current use									
7. Any method (%)	47.2	65.7	40.5	44.1	54.9	49.7	62.5	40.3	31.9
8. Any modern method (%)	44.4	62.0	38.0	41.6	52.1	46.5	57.7	38.1	31.0
8a. Female sterilization (%)	34.2	39.6	32.2	36.0	39.5	30.7	24.1	30.8	25.4
8b. Male sterilization (%)	0.8	1.7	0.5	0.9	0.6	0.8	0.6	1.5	2.4
8c. IUD (%)	1.6	2.4	1.3	1.2	2.1	1.3	4.9	1.2	1.2
8d. Pill (%)	2.0	4.9	1.0	0.8	1.5	4.8	5.5	1.5	0.5
8e. Condom (%)	5.7	13.0	3.1	2.7	8.4	8.6	21.9	3.1	1.5
Unmet need for family planning									
9. Total unmet need (%)	14.6	9.8	16.3	14.2	9.7	17.5	14.0	17.6	19.8
9a. For spacing (%)	7.3	5.2	8.0	5.9	4.1	13.1	8.1	8.7	10.8
9b. For limiting (%)	7.3	4.6	8.3	8.4	5.6	4.4	5.9	8.9	9.0
Maternal and Child Health									
Maternity care (for births in the last 3 years)									
10. Mothers who had at least 3 antenatal care visits for their last birth (%)	41.2	74.7	32.5	28.5	(38.5)	66.5	86.0	23.6	18.1
11. Mothers who consumed IFA for 90 days or more when they were pregnant with their last child (%)	12.8	30.7	8.1	6.2	(16.6)	17.0	56.9	na	na
12. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%) ²	43.2	77.0	34.6	31.2	52.3	64.1	90.3	35.8	19.3
13. Institutional births (%) ²	32.2	67.7	23.3	20.6	35.1	53.6	78.9	21.5	12.0
14. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/other health personnel within 2 days of delivery for their last birth (%) ²	29.0	56.8	21.9	19.7	(28.2)	42.4	75.5	na	na
Child immunization and vitamin A supplementation²									
15a. Children 12-23 months fully immunized (BCG, measles, and 3 doses each of polio/DPT) (%)	26.5	44.3	22.1	20.5	*	38.5	(61.4)	17.3	21.1
15b. Children 12-23 months who have received BCG (%)	68.5	76.1	66.7	61.6	*	80.4	(95.8)	53.9	45.6
15c. Children 12-23 months who have received 3 doses of polio vaccine (%)	65.2	73.9	63.0	58.9	*	79.2	(74.4)	44.6	33.9
15d. Children 12-23 months who have received 3 doses of DPT vaccine (%)	38.7	63.6	32.6	30.8	*	52.4	(88.3)	26.1	29.6
15e. Children 12-23 months who have received measles vaccine (%)	42.7	60.2	38.4	34.3	*	55.5	(86.0)	27.1	31.3
16. Children age 12-35 months who received a vitamin A dose in last 6 months (%)	16.5	23.4	14.7	11.7	(9.4)	23.5	44.2	na	na
Treatment of childhood diseases (children under 3 years)²									
17. Children with diarrhoea in the last 2 weeks who received ORS (%)	16.0	28.0	11.5	11.7	*	(18.6)	*	20.3	12.1
18. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	56.6	64.0	53.8	52.8	*	(57.5)	*	58.2	50.0
19. Children with acute respiratory infection or fever in the last 2 weeks taken to a health facility (%)	68.9	80.9	65.7	69.2	*	(78.1)	*	na	na
Child Feeding Practices and Nutritional Status of Children^{2, 3}									
20. Children under 3 years breastfed within one hour of birth (%)	13.3	21.0	11.3	10.5	(10.8)	18.4	25.3	4.8	8.1
21. Children age 0-5 months exclusively breastfed (%)	33.2	21.2	36.2	34.4	*	20.2	*	na	na
22. Children age 6-9 months receiving solid or semi-solid food and breastmilk (%)	38.7	60.7	34.0	32.3	*	(44.6)	*	na	na
23. Children under 3 years who are stunted (%)	40.1	29.4	42.8	45.6	(36.7)	32.1	16.3	59.0	45.5
24. Children under 3 years who are wasted (%)	22.5	19.6	23.3	24.7	(31.8)	17.6	12.6	16.2	23.8
25. Children under 3 years who are underweight (%)	36.8	26.1	39.5	43.1	(35.9)	24.6	17.1	46.7	41.8
Nutritional Status of Ever-Married Adults (age 15-49)									
26. Women whose Body Mass Index is below normal (%)	33.6	25.6	36.5	35.5	33.8	33.3	18.9	36.1	na
27. Men whose Body Mass Index is below normal (%)	33.8	22.8	38.5	39.1	45.7	36.9	18.9	na	na
28. Women who are overweight or obese (%)	10.2	22.1	5.8	7.4	12.2	13.1	25.4	7.1	na
29. Men who are overweight or obese (%)	8.4	19.3	3.6	3.0	3.8	6.5	18.7	na	na
Anaemia among Children and Adults									
30. Children age 6-35 months who are anaemic (%)	79.1	78.5	80.2	80.5	77.5	81.3	72.3	82.3	na
31. Ever-married women age 15-49 who are anaemic (%)	53.8	48.5	55.7	55.7	48.8	57.1	34.9	48.5	na
32. Pregnant women age 15-49 who are anaemic (%)	61.7	64.5	60.8	61.3	*	64.5	*	51.4	na
33. Ever-married men age 15-49 who are anaemic (%)	21.2	16.7	23.1	29.1	32.1	19.7	10.9	na	na
Knowledge of HIV/AIDS among Ever-Married Adults (age 15-49)									
34. Women who have heard of AIDS (%)	33.8	73.0	19.3	14.8	39.7	73.1	98.8	20.8	na
35. Men who have heard of AIDS (%)	74.2	96.5	64.5	41.7	56.1	86.3	98.7	na	na
36. Women who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	27.3	61.6	14.7	9.8	31.1	59.9	95.5	na	na
37. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	63.2	88.6	52.1	26.0	44.5	73.9	95.2	na	na
Women's Empowerment									
38. Currently married women who usually participate in household decisions (%)	22.8	37.9	17.3	20.5	20.0	22.8	42.2	na	na
39. Ever-married women who have ever experienced spousal violence (%)	46.3	45.4	46.6	50.4	54.1	42.2	20.4	na	na

na : Not available

() Based on 25-49 unweighted cases

* Based on fewer than 25 unweighted cases

¹ Excludes pregnant women.

² Based on the last 2 births in the 3 years before the survey to ever-married women.

³ Based on WHO standard.

⁴ For children education refers to mother's education. Children with missing information on the mother's education are not included in the education columns.