

Key Indicators for Uttarakhand from NFHS-3	NFHS-3 (2005-06)	Residence		Education				NFHS-2 (1998-99)	NFHS-1 (1992-93)
		Urban	Rural	No education ⁴	<5 years complete ⁴	5-9 years complete ⁴	10 years complete and above ⁴		
Marriage and Fertility									
1. Women age 20-24 married by age 18 (%)	23.0	16.3	25.2	51.0	*	30.7	4.2	25.9	na
2. Men age 25-29 married by age 21 (%)	21.3	7.1	30.0	*	*	37.9	8.1	na	na
3. Total fertility rate (children per woman)	2.6	2.2	2.7	3.6	2.5	2.5	2.0	2.6	na
4. Women age 15-19 who were already mothers or pregnant at the time of the survey (%)	6.2	1.7	7.8	26.5	*	5.8	0.9	na	na
5. Median age at first birth for women age 25-49	20.5	21.5	20.3	19.3	19.4	20.1	23.3	20.1	na
6. Married women with 2 living children wanting no more children ¹ (%)	87.3	87.9	86.9	78.0	*	88.8	91.3	71.8	na
6a. Two sons	93.0	93.0	93.0	86.6	*	96.5	96.2	93.2	na
6b. One son, one daughter	91.3	91.7	91.1	86.7	*	87.1	95.4	70.2	na
6c. Two daughters	49.4	(60.7)	(40.6)	*	nc	*	(62.6)	35.4	na
Family Planning (currently married women, age 15-49)									
Current use									
7. Any method (%)	59.3	65.3	57.2	58.4	69.0	55.8	62.7	43.1	na
8. Any modern method (%)	55.5	59.2	54.2	55.1	61.2	53.1	57.5	40.4	na
8a. Female sterilization (%)	32.1	20.4	36.3	41.9	45.8	33.8	14.4	27.3	na
8b. Male sterilization (%)	1.8	2.7	1.4	1.8	4.9	1.6	1.3	3.8	na
8c. IUD (%)	1.5	2.7	1.0	0.5	1.3	1.9	2.6	1.6	na
8d. Pill (%)	4.2	4.7	4.0	2.8	3.7	4.1	6.3	1.4	na
8e. Condom (%)	15.7	28.3	11.2	8.1	5.4	11.7	32.1	6.2	na
Unmet need for family planning									
9. Total unmet need (%)	10.8	9.8	11.2	8.2	7.8	16.0	10.3	21.0	na
9a. For spacing (%)	4.4	3.2	4.8	2.9	2.7	6.9	4.4	10.5	na
9b. For limiting (%)	6.5	6.6	6.4	5.3	5.1	9.1	5.9	10.5	na
Maternal and Child Health									
Maternity care (for births in the last 3 years)									
10. Mothers who had at least 3 antenatal care visits for their last birth (%)	44.8	71.5	36.2	21.2	*	39.0	80.9	19.7	na
11. Mothers who consumed IFA for 90 days or more when they were pregnant with their last child (%)	26.2	43.0	20.8	8.9	*	18.8	55.5	na	na
12. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%) ²	41.6	64.6	34.6	21.5	*	34.1	76.7	34.6	na
13. Institutional births (%) ²	36.1	59.9	28.8	15.6	*	29.1	71.7	20.6	na
14. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/other health personnel within 2 days of delivery for their last birth (%) ²	30.2	52.3	23.1	11.8	*	20.7	64.2	na	na
Child immunization and vitamin A supplementation²									
15a. Children 12-23 months fully immunized (BCG, measles, and 3 doses each of polio/DPT) (%)	60.0	67.2	57.4	40.4	*	59.6	86.4	40.9	na
15b. Children 12-23 months who have received BCG (%)	83.5	84.4	83.2	66.6	*	91.1	98.5	76.8	na
15c. Children 12-23 months who have received 3 doses of polio vaccine (%)	80.3	82.8	79.4	63.9	*	86.6	95.5	62.4	na
15d. Children 12-23 months who have received 3 doses of DPT vaccine (%)	67.1	68.7	66.5	45.5	*	67.3	95.5	56.1	na
15e. Children 12-23 months who have received measles vaccine (%)	71.6	75.0	70.3	56.6	*	68.5	94.0	56.0	na
16. Children age 12-35 months who received a vitamin A dose in last 6 months (%)	20.3	23.6	19.1	11.9	*	18.5	33.5	na	na
Treatment of childhood diseases (children under 3 years)²									
17. Children with diarrhoea in the last 2 weeks who received ORS (%)	35.6	(45.2)	32.6	(21.7)	*	(34.8)	(60.8)	31.5	na
18. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	64.8	(71.0)	62.8	(64.0)	*	(68.8)	(66.8)	64.0	na
19. Children with acute respiratory infection or fever in the last 2 weeks taken to a health facility (%)	71.6	(80.0)	69.5	71.5	*	(61.9)	(82.1)	na	na
Child Feeding Practices and Nutritional Status of Children^{2,3}									
20. Children under 3 years breastfed within one hour of birth (%)	32.9	29.1	34.2	33.9	*	29.6	34.9	24.1	na
21. Children age 0-5 months exclusively breastfed (%)	31.2	*	35.1	(36.4)	*	(28.5)	(27.6)	na	na
22. Children age 6-9 months receiving solid or semi-solid food and breastmilk (%)	51.6	*	51.4	(39.4)	*	(56.9)	(54.5)	na	na
23. Children under 3 years who are stunted (%)	39.6	22.1	44.9	48.2	*	41.2	24.9	52.5	na
24. Children under 3 years who are wasted (%)	18.2	9.8	21.0	24.3	*	18.4	10.9	9.0	na
25. Children under 3 years who are underweight (%)	31.7	20.9	35.2	43.0	*	34.2	13.5	36.3	na
Nutritional Status of Ever-Married Adults (age 15-49)									
26. Women whose Body Mass Index is below normal (%)	25.7	11.1	30.8	32.8	30.8	26.9	13.0	32.4	na
27. Men whose Body Mass Index is below normal (%)	21.8	13.2	25.6	35.3	*	27.4	9.9	na	na
28. Women who are overweight or obese (%)	16.0	30.5	11.0	8.2	21.0	14.0	29.3	9.2	na
29. Men who are overweight or obese (%)	11.4	18.6	8.2	7.3	*	6.3	18.3	na	na
Anaemia among Children and Adults									
30. Children age 6-35 months who are anaemic (%)	69.2	66.2	70.8	74.4	*	71.8	60.1	77.4	na
31. Ever-married women age 15-49 who are anaemic (%)	54.7	49.3	56.4	57.6	56.0	54.7	49.6	45.6	na
32. Pregnant women age 15-49 who are anaemic (%)	50.8	(45.7)	52.3	54.6	*	(54.3)	(41.5)	49.8	na
33. Ever-married men age 15-49 who are anaemic (%)	30.5	23.2	33.7	42.0	*	36.5	19.0	na	na
Knowledge of HIV/AIDS among Ever-Married Adults (age 15-49)									
34. Women who have heard of AIDS (%)	64.3	84.5	57.2	37.2	66.4	71.1	98.3	35.6	na
35. Men who have heard of AIDS (%)	91.0	93.7	89.7	62.5	*	93.6	100.0	na	na
36. Women who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	52.2	71.7	45.3	25.6	44.1	56.1	89.5	na	na
37. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	81.4	84.8	79.6	42.0	*	84.0	95.8	na	na
Women's Empowerment									
38. Currently married women who usually participate in household decisions (%)	36.0	46.7	32.2	33.3	35.1	33.3	42.7	na	na
39. Ever-married women who have ever experienced spousal violence (%)	27.8	22.8	29.6	39.4	30.6	28.8	8.6	na	na

na : Not available

nc : Not calculated because there are no cases

() Based on 25-49 unweighted cases

* Based on fewer than 25 unweighted cases

¹ Excludes pregnant women.

² Based on the last 2 births in the 3 years before the survey to ever-married women.

³ Based on WHO standard.

⁴ For children education refers to mother's education. Children with missing information on the mother's education are not included in the education columns.